

## WHAT IS CLAIMED IS:

1. A method of preventing and/or treating diabetes type 2 in a subject in need thereof, said method comprising step of administering pharmaceutically effective amount of an extract of plant *Pueraria tuberosa* or butanol fraction of the extract or Lupinoside A4 (LPA<sub>4</sub>), optionally along with additive(s) to the subject.
2. A method as claimed in claim 1, wherein the subject is an animal.
3. A method as claimed in claim 1, wherein the subject is a human being.
4. A method as claimed in claim 1, wherein the extract is obtained from root of the plant.
5. A method as claimed in claim 1, wherein the additive is selected from a group comprising nutrients such as proteins, carbohydrates, sugars, talc, magnesium stearate, cellulose, calcium carbonate, starch, gelatin paste, pharmaceutically acceptable carrier, excipients, diluent and, solvent.
6. A method as claimed in claim 1, wherein the fraction is administered at the concentration ranging between 1 to 40 mg /kg body weight.
7. A method as claimed in claim 1, wherein the Lupinoside is administered at the concentration ranging between 1 to 40 mg /kg body weight.
8. A method as claimed in claim 1, wherein the administration route is selected from a group comprising orally, intravenously, intramuscularly, and subcutaneously.
9. A pharmaceutical composition useful in preventing and/or treating diabetes type 2, said composition comprising an extract of plant *Pueraria tuberosa* or butanol fraction of the extract or Lupinoside A4 (LPA<sub>4</sub>), and additive(s).
10. A pharmaceutical composition as claimed in claim 9, wherein the additive is selected from a group comprising nutrients such as proteins, carbohydrates,

sugars, talc, magnesium stearate, cellulose, calcium carbonate, starch, gelatin paste, pharmaceutically acceptable carrier, excipients, diluent and, solvent.

5 11. A pharmaceutical composition as claimed in claim 9, the extract is obtained from root of the plant.

12. A pharmaceutical composition as claimed in claim 9, the fraction is of concentration ranging between 1 to 40 mg /kg body weight.

10 13. A pharmaceutical composition as claimed in claim 9, the Lupinoside is of concentration ranging between 1 to 40 mg /kg body weight.

14. A pharmaceutical composition as claimed in claim 9, wherein the composition is in a form selected from a group comprising capsule, syrup, concentrate, powder, 15 and granules.

15. A pharmaceutical composition as claimed in claim 9, wherein the extract is an aqueous extract.

20 16. A method of augmenting Glut4 phosphorylation and Glut4 translocation to a target cell membrane to enhance insulin signal in a signal transduction pathway in a subject in need thereof, said method comprising administering pharmaceutically effective amount of an extract of plant *Pureria tuberosa* or butanol fraction of the extract or Lupinoside A4 (LPA<sub>4</sub>), optionally along with 25 additive(s) to the subject.

17. A method as claimed in claim 16, wherein the subject is an animal.

18. A method as claimed in claim 16, wherein the subject is a human being. 30

19. A method as claimed in claim 16, wherein the extract is obtained from root of the plant.

- 5
20. A method as claimed in claim 16, wherein the additive is selected from a group comprising nutrients such as proteins, carbohydrates, sugars, talc, magnesium stearate, cellulose, calcium carbonate, starch, gelatin paste, pharmaceutically acceptable carrier, excipients, diluent and, solvent.
21. A method as claimed in claim 16, wherein the fraction is administered at the concentration ranging between 1 to 40 mg /kg body weight.
- 10 22. A method as claimed in claim 16, wherein the Lupinoside is administered at the concentration ranging between 1 to 40 mg /kg body weight.
23. A method as claimed in claim 16, wherein the method helps prevent/treat type 2 diabetes.
- 15 24. A method as claimed in claim 16, wherein the method shows increase in glucose uptake by the cells.
25. A method as claimed in claim 16, wherein the method is non-toxic to the cells.
- 20 26. A method as claimed in claim 16, wherein the translocation is from cytosol to membrane of the insulin response cells.
27. A method as claimed in claim 16, wherein the Lupinoside A<sub>4</sub> (LP<sub>4</sub>) prevents palmitate induced defects on insulin signaling.
- 25 28. A method as claimed in claim 16, wherein the Lupinoside A<sub>4</sub> (LP<sub>4</sub>) allows insulin to stimulate IR-beta and Akt phosphorylation.
- 30 29. A simplified and inexpensive process of obtaining extract and thereafter selectively, its active n-butanol fraction and active molecule Lupinoside PA (LPA<sub>4</sub>), useful in preventing and/or treating diabetes type 2, said process comprising steps of:
- a. cutting the plant parts into small parts,

- 5
- b. extracting the cut parts with methanol and water,
  - c. partitioning the methanol and water extract between ethyl acetate and water,
  - d. extracting the aqueous layer further with n-butanol to obtain butanol fraction, and
  - 10 e. subjecting the n-butanol fraction to chromatography with water and methanol as eluent to obtain Lupinoside PA<sub>4</sub> (LPA<sub>4</sub>).

30. A method as claimed in claim 29, wherein the plant part is root.

15 31. A method as claimed in claim 29, wherein the solvent is selected from a group comprising methanol, and water.

32. A method as claimed in claim 29, wherein the water and methanol are in the ratio of about 1:1.

20 33. A method as claimed in claim 29, wherein the chromatography is column chromatography.

25